



### What is Mytesi?

Mytesi is a prescription medicine that helps relieve symptoms of diarrhea not caused by an infection in adults living with HIV/AIDS on antiretroviral therapy (ART).

# HOW TO TAKE MYTESI

1 Mytesi 125 mg DELAYED-RELEASE TABLET IS TAKEN **2 TIMES EACH DAY**

YOU CAN TAKE IT WITH OR WITHOUT FOOD  / 

Mytesi tablets should **NOT** be crushed or chewed

Tablets should be **swallowed whole** 

NapoCares™ Patient Support Programs can help you start and continue your Mytesi® treatment.

To enroll in NapoCares or to learn more about its programs, please call (888) 527-NAPO (6276) Monday-Friday 8 am-5 pm CST.

*All NapoCares programs are free of charge for Mytesi patients.*



**Copay Savings Program** to help you save on your out-of-pocket costs<sup>†</sup>



**Patient Assistance Program** for uninsured patients



Support to help you **understand your insurance coverage**



Support for **Prior Authorizations and Appeals**



**Mail-order** pharmacy support



**Patient call center** to answer any questions



**Linkage to specialty pharmacies** to fill your prescription

### Important Safety Information

Mytesi (crofelemer) is not approved to treat infectious diarrhea (diarrhea caused by bacteria, a virus, or a parasite). Before starting you on Mytesi, your healthcare provider will first be sure that you do not have infectious diarrhea. Otherwise, there is a risk you would not receive the right medicine and your infection could get worse. In clinical studies, the most common side effects that occurred more often than with placebo were upper respiratory tract (sinus, nose, and throat) infection (6%), bronchitis (4%), cough (4%), flatulence (3%), and increased bilirubin (3%).

See package insert for full prescribing information.

<sup>\*</sup>Subject to eligibility. Restrictions apply. Rx only. Subject to change or cancellation at any time and without notice.

<sup>†</sup>Patient must be commercially insured for eligibility. Subject to change or cancellation at any time and without notice.

# TRACKING YOUR PROGRESS

When taking Mytesi daily, most patients continue to see improvement in their symptoms even up to 20 weeks after starting treatment. It's important to continue taking Mytesi daily to see these improvements, even if the symptom relief seems gradual.


## WHAT'S YOUR NUMBER?

Adapted from the Bristol Stool Form Scale


### CONSTIPATION



### NORMAL



### DIARRHEA (LOOSE/WATERY STOOLS)



Date	Number of Loose/Watery Stools (5, 6, or 7 from above)	Notes
<b>BASELINE</b> before starting Mytesi Date: ___/___/___		
<b>1 WEEK</b> after starting Mytesi Date: ___/___/___		
<b>4 WEEKS</b> after starting Mytesi Date: ___/___/___		
<b>8 WEEKS</b> after starting Mytesi Date: ___/___/___		
<b>20 WEEKS</b> after starting Mytesi Date: ___/___/___		

NapoCares has support for Mytesi patients at no cost to them, whether they are uninsured, underinsured, or insured. To enroll in NapoCares or to learn more about its programs, please call 1-888-527-NAPO (6276) Monday-Friday 8 am - 5 pm CST.

Important Safety Information on the other side. See package insert for full prescribing information.